

EXPOSURE TO COVID-19 TRIAGE CHART

DISCLAIMER: THIS CHART IS TO ASSIST IN COUNSELING AND IS NOT DEFINITIVE

Scenario	Risk Factor	Nasopharyngeal Swab	Antibody Test	Quarantine
Briefly walking by a person who tested positive for COVID-19 and was not experiencing symptoms.	MINIMAL RISK	LOW VALUE	LOW VALUE	NOT NECESSARY Highly suggest to wear mask around others.
Briefly being in the same room AND within 6 feet of a person who tested positive for COVID-19, 48 hours before they started with symptoms or had symptoms.	LOW RISK	MEDIUM VALUE http://bluefishmd.com/wp-content/uploads/2020/06/6.22.2020-COVID-19-Testing-FacilitiesC.pdf	LOW VALUE	NOT NECESSARY Mandatory mask wearing around others for 14 days minimum.
Prolonged close contact (15 minutes or longer) AND within 6 feet of a person who tested positive for COVID-19, 48 hours before they started with symptoms or had symptoms.	MEDIUM RISK	HIGH VALUE http://bluefishmd.com/wp-content/uploads/2020/06/6.22.2020-COVID-19-Testing-FacilitiesC.pdf	MEDIUM VALUE https://bluefishmd.com/wp-content/uploads/2020/05/5.20.20.-Updated-COVID-Antibody-Handout.pdf	STRONGLY CONSIDER Mandatory mask wearing around others for 14 days minimum.
Sustained close contact within the same room at work, school, or home with person who tested positive for COVID-19.	HIGH RISK	HIGH VALUE http://bluefishmd.com/wp-content/uploads/2020/06/6.22.2020-COVID-19-Testing-FacilitiesC.pdf	MEDIUM VALUE https://bluefishmd.com/wp-content/uploads/2020/05/5.20.20.-Updated-COVID-Antibody-Handout.pdf	MANDATORY Should not go to work or school/daycare or be around others for 14 days minimum.* Other family members need only quarantine if the original quarantined individual becomes symptomatic or has a positive test.

Please note if both parties were wearing masks it significantly reduces the risk of transmission in all scenarios.

* https://bluefishmd.com/wp-content/uploads/2020/07/Home-IsolationQuarantine-Release-Graphic_FINAL_2nd_Chart.pdf