

EXPOSURE TO COVID-19 TRIAGE CHART

Scenario	Risk Factor	PCR Test	Antigen Test	Quarantine
Briefly walking by a person who tested positive for COVID-19 and was not experiencing symptoms.	MINIMAL RISK	LOW VALUE	LOW VALUE	NOT NECESSARY Highly suggest to wear mask around others
In the same room for less than 15 minutes AND within 6 feet of a person who tested positive for COVID-19, 48 hours before the test of onset of symptoms.	LOW RISK	MEDIUM VALUE https://bluefishmd.com/wp-content/uploads/2020/09/8.28.2020-COVID-19-Testing-Facilities-for-Website.pdf	LOW VALUE	NOT NECESSARY Mandatory mask wearing around others for 14 days minimum
In the same room within 6 feet for a cumulative of 15 minutes or longer over a 24 hours period, 48 hours before the close contact started symptoms or had symptoms.	HIGH RISK	HIGH VALUE https://bluefishmd.com/wp-content/uploads/2020/09/8.28.2020-COVID-19-Testing-Facilities-for-Website.pdf	MEDIUM VALUE	MANDATORY FOR NON HEALTHCARE WORKERS
Sustained close contact for a cumulative of 15 minutes or longer within the same room at work, school, or home with person who tested positive for COVID-19.	HIGHEST RISK	HIGH VALUE https://bluefishmd.com/wp-content/uploads/2020/09/8.28.2020-COVID-19-Testing-Facilities-for-Website.pdf	MEDIUM VALUE	MANDATORY Should not go to work or school/daycare or e around others for 14 days minimum. * Other family members need only quarantine if the original quarantined individual becomes symptomatic or has a positive test.

Please note if both parties were wearing masks it is likely to significantly reduce the risk of transmission in all scenarios.

*https://bluefishmd.com/wp-content/uploads/2020/07/Home-IsolationQuarantine-Release-Graphic_FINAL_2nd_Chart.pdf