

# BLUE FISH

P E D I A T R I C S  
www.bluefishmd.com

Patient Name: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Excerpt from the CDC website:

<https://www.cdc.gov/vaccines/covid-19/info-by-product/clinical-considerations.html>

Co-administration with other vaccines

None of the currently authorized COVID-19 vaccines are live virus vaccines. Because data are lacking on the safety and efficacy of COVID-19 vaccines administered simultaneously with other vaccines, **the vaccine series should routinely be administered alone, with a minimum interval of 14 days before or after administration of any other vaccine.** However, COVID-19 and other vaccines may be administered within a shorter period in situations where the benefits of vaccination are deemed to outweigh the potential unknown risks of vaccine coadministration (e.g., tetanus-toxoid-containing vaccination as part of wound management, rabies vaccination for post-exposure prophylaxis, measles or hepatitis A vaccination during an outbreak) or to avoid barriers to or delays in to COVID-19 vaccination (e.g., in long-term care facility residents or healthcare personnel who received influenza or other vaccinations before or upon admission or onboarding). If COVID-19 vaccines are administered within 14 days of another vaccine, doses do not need to be repeated for either vaccine.

**My child has not received the COVID-19 vaccine in the past 14 days. I also acknowledge that my child should wait 14 days after receiving any other type of vaccines before receiving the COVID-19 vaccine.**

Parent/Legal Guardian Name: \_\_\_\_\_

Parent/Legal Guardian Signature: \_\_\_\_\_ Date: \_\_\_\_\_